



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

II. Solution by E. B. ESCOTT, Ann Arbor, Michigan.

The solution of the congruence $x^2 + y^2 \equiv 0 \pmod{5}$, is $x \equiv \pm 2y \pmod{5}$.
Let $x = 5a \pm 2y$, and substitute in $x^2 + y^2 \equiv 0 \pmod{5^2}$.

We get $x \equiv \pm 7y \pmod{5^2}$, and proceeding in this way we get finally,
 $x \equiv \pm 182y \pmod{5^4}$.

NOTE. A similar and more complete solution by the Proposer has already been published. ED. W.

180. Proposed by A. H. HOLMES, Brunswick, Maine.

Find integral values of x and y such that $96x - 96y + 21 = \square$.

II. Solution by B. KRAMER, E. M., Pittsburg, Pennsylvania.

Let $x - y = z$. Then $96z + 21 = u^2$. From this it is easily seen that u is odd, and $u^2 \equiv 5 \pmod{8}$.

But if we put $u = 2k + 1$, $u^2 = 4k(k + 1) + 1$, which, since k or $k + 1$ is even, says that $u^2 \equiv 1 \pmod{8}$. Hence no square can be congruent to 5 modulo 8, and the relation is impossible.

181. Proposed by V. M. SPUNAR, M. and E. E., Chicago, Ill.

If $2n + 1$ is an odd prime p , $(2n)! \equiv (-1)^{n2^n} (n!)^2 \pmod{p^2}$.

Solution by S. LEFSCHETZ, Clark University.

The given congruence can be written

$$\begin{aligned} 2n! &\equiv (-1)^{n2^{2n}} [2 \cdot 4 \dots 2n]^2 \pmod{p^2}, \text{ or} \\ 1 \cdot 3 \cdot 5 \dots (2n-1) &\equiv (-1)^{n2^{2n}} \cdot 2 \cdot 4 \cdot 6 \dots 2n \pmod{p^2} \\ &\equiv (-1)^{n2^{2n}} (p-1)(p-3) \dots [p-(2n-1)] \\ &\equiv 2^{2n} [1 \cdot 3 \dots (2n-1) - p \cdot 1 \cdot 2 \dots (2n-1) \left(\frac{1}{1} + \frac{1}{3} + \dots + \frac{1}{2n-1}\right)] \pmod{p^2}. \end{aligned}$$

If we adopt Bachmann's notation [*Niedere Zahlentheorie*, Bd. 1, p. 161] and call $1/m$ the number m' such that $mm' \equiv 1 \pmod{p}$. We have to prove, as can be seen at once that:

$$\frac{2^{2n}-1}{p} \equiv \frac{1}{1} + \frac{1}{3} + \dots + \frac{1}{2n-1} \pmod{p},$$

and this is proved by Bachmann (loc. cit., p. 164).

182. Proposed by V. M. SPUNAR, M. and E. E., Chicago, Ill.

Find two general solutions in integers of the equation $x^2 = 616318177y - 1$.

No solution of this problem has been received.